



Kyphoplasty/Vertebroplasty Post-operative Instructions

Successful surgery depends not only on the success of the procedure, but also on your cooperation and compliance during the post-operative period. To assist you in this process, below are instructions and guidelines to follow during your recovery period.

Wound

- There will be a small incision on your back. Your incision will be assessed at your post-operative office visit.
- Begin changing your dressing daily after discharge from the hospital unless directed otherwise by someone from our clinical staff.
- You may shower but you need to cover the incision with a plastic covering to keep it dry. **Do not soak the incision in water for 6 weeks.**
- Do not put creams or ointments on the wound for the first 3 weeks – or until incision is healed well.

Pain Relief

- Our office will provide you with prescription pain medication to help with your surgical pain. You should use these medications sparingly and only as directed. It is expected that most of your pain can be managed with over the counter medications such as Tylenol.
- As your incision heals, we would expect you to begin weaning from prescription medications. This should begin about a week after surgery.

- If you are prescribed antibiotics after surgery, please finish them in their entirety.
- You will need to avoid anti-inflammatory medications for 1 week before and 2 weeks following surgery.
- You will need to avoid blood thinners (Aspirin, Coumadin, Lovenox, Plavix) for 1 week prior to surgery. In most cases, you may restart them (at the doctor's discretion) 4 days after surgery or 4 days after drains are removed.
- **Remember, ABSOLUTELY NO MEDICATION REFILLS WILL BE PROCESSED ON THE WEEKEND.** Please plan accordingly. Contact your pharmacy to process all refill requests.

Exercise/Activity

- Avoid riding in a car for 2 weeks – unless medically necessary.
- The best exercise is walking. Small amounts done frequently are best. Try to set a goal to walk a little more each day. This of course will depend on your individual situation and abilities.
- Wear the white compression stockings given to you at the hospital for 2 weeks following surgery. You can take them off when walking but should wear them at night and when inactive for prolonged periods
- **DO NOT SMOKE.** Smoking may delay the healing of your surgical wound. See your primary care physician if you need assistance to quit smoking.

Driving

- You will not be able to drive for approximately 2 weeks after this type of surgery.
- It is the policy of this office to advise you not to drive while under the influence of pain medications.

Return to Work

- If you can work from home, you can begin doing light desk/phone work as tolerated about 1-2 weeks after surgery.
- If you have to report to the office to perform your job, you will be able to do this at 2-3 weeks after surgery. You will be restricted to light desk/phone type work.
- If your job requires heavy physical activity, you will be unable to perform this type of work for at least 12 weeks after surgery.

Follow-up

- You have a follow-up scheduled with one of our clinical staff in our Plano office on _____ . Please arrive at least 10 minutes early to check in.

What to Watch For

- Please contact our office for any of the following:
Oral temperature greater than 100.5 ° F
Excessive redness, swelling, or drainage at the incision site
- For general medical problems such as sore throat, cough, nasal drainage, etc., please contact your primary care physician. You should also contact your primary care physician regarding in questions about your routine medications not prescribed by this office.
- For medical emergencies, please call 911 or report to the nearest emergency room.



6160 Windhaven Pkwy, Suite 200 Plano, TX 75093
Ph. (972)378-6908 Fax (972) 378-6586

Date of Surgery: _____ **Scheduled Surgery Time:** _____ **Hospital Arrival Time:** _____ **No food or drink after midnight day prior to surgery.**